

# CONSTRAINED OPTIMIZATION

A general constrained optimization problem has the form

$$\begin{aligned} \min_x & f(x) \\ \text{s.t.} & g_i(x) \leq 0, \quad i=1, \dots, m \\ & h_i(x) = 0, \quad i=1, \dots, p \end{aligned}$$

where  $x \in \mathbb{R}^d$ . If  $x$  satisfies all the constraints,

Ⓐ we say  $x$  is \_\_\_\_\_. Assume  $f$  is defined on all feasible points.

## Lagrangian Duality

The \_\_\_\_\_ function is


$$L(x, \lambda, \nu) := f(x) + \sum_{i=1}^m \lambda_i g_i(x) + \sum_{i=1}^p \nu_i h_i(x)$$

and  $\lambda = [\lambda_1, \dots, \lambda_m]^T$  and  $\nu = [\nu_1, \dots, \nu_p]^T$  are called \_\_\_\_\_ or \_\_\_\_\_.

The (Lagrange) dual function is

$$L_D(\lambda, \nu) := \min_x L(x, \lambda, \nu)$$

Note  $L_D$  is concave, being the point-wise minimum of a family of affine functions



③ Then we can define the                      optimization problem:

$$\max_{\lambda, \nu: \lambda_i \geq 0} L_D(\lambda, \nu)$$

Why would we constrain  $\lambda_i \geq 0$ ?

### The Primal

Similarly, we could define the                      function

$$L_P(x) := \max_{\lambda, \nu: \lambda_i \geq 0} L(x, \lambda, \nu)$$

and the primal optimization problem

$$\min_x L_P(x) = \min_x \max_{\lambda, \nu: \lambda_i \geq 0} L(x, \lambda, \nu)$$

It is like the dual but the min + max are swapped.

Notice that

$$L_p(x) = \begin{cases} f(x) & \text{if } x \text{ is feasible} \\ -\infty & \text{else} \end{cases}$$

Thus the primal encompasses the original problem (i.e., they have the same solution).

### Weak Duality

Claim  $d^* := \max_{\lambda, \nu: \lambda_i \geq 0} \min_x L(x, \lambda, \nu)$

$$\leq \min_x \max_{\lambda, \nu: \lambda_i \geq 0} L(x, \lambda, \nu) =: p^*$$

Proof: Let  $\tilde{x}$  be feasible. Then for any  $\lambda, \nu$  with  $\lambda_i \geq 0$ ,

$$L(\tilde{x}, \lambda, \nu) = f(\tilde{x}) + \sum \lambda_i g_i(\tilde{x}) + \sum \nu_i h_i(\tilde{x}) \leq f(\tilde{x})$$

Hence

$$L_D(\lambda, \nu) = \min_x L(x, \lambda, \nu) \leq L(\tilde{x}, \lambda, \nu) \leq f(\tilde{x}).$$

This is true for any feasible  $\tilde{x}$ , so

$$L_D(\lambda, \nu) \leq \min_{\substack{\tilde{x} \\ \tilde{x} \text{ feasible}}} f(\tilde{x}) = p^*$$

Taking the max over  $\lambda, \nu: \lambda_i \geq 0$ , we have

$$d^* = \max_{\lambda, \nu: \lambda_i \geq 0} L_D(\lambda, \nu) \leq p^*$$

□

© The difference  $p^* - d^*$  is called the \_\_\_\_\_.

### Strong Duality

If  $p^* = d^*$ , we say strong duality holds.

Theorem | If the primal problem is \_\_\_\_\_ ( $f, g_i$  convex,  $h_i$  affine), and a constraint qualification holds, then  $p^* = d^*$ .

Examples | of constraint qualifications:

- All  $g_i$  are \_\_\_\_\_.
- $\exists x$  s.t.  $h_i(x) = 0 \ \forall i$ ,  $g_i(x) < 0 \ \forall i$   
(strict feasibility)

### KKT Conditions

Assume  $f, g_i, h_i$  are differentiable.

### Necessity

Theorem | If  $p^* = d^*$ ,  $x^*$  is primal optimal, and  $(\lambda^*, \nu^*)$  is dual optimal, then the

Karush-Kuhn-Tucker conditions hold:

KKT conditions

$$\left[ \begin{array}{l} (1) \nabla f(x^*) + \sum_{i=1}^m \lambda_i^* \nabla g_i(x^*) + \sum_{i=1}^p \nu_i^* \nabla h_i(x^*) = 0 \\ (2) g_i(x) \leq 0, \quad i=1, \dots, m \\ (3) h_i(x^*) = 0, \quad i=1, \dots, p \\ (4) \lambda_i^* \geq 0, \quad i=1, \dots, m \\ (5) \lambda_i^* g_i(x^*) = 0, \quad i=1, \dots, m \quad (\text{complementary slackness}) \end{array} \right.$$

Proof: To prove the first condition, notice that  $L(x, \lambda^*, \nu^*)$  is minimized by  $x^*$ , and thus its gradient wrt  $x$  must be 0 at  $x^*$ .

To prove the last condition, write

$$\begin{aligned} f(x^*) &= L_D(\lambda^*, \nu^*) \\ &= \min_x \left( f(x) + \sum_{i=1}^m \lambda_i^* g_i(x) + \sum_{i=1}^p \nu_i^* h_i(x) \right) \\ &\leq f(x^*) + \sum_{i=1}^m \lambda_i^* g_i(x^*) + \sum_{i=1}^p \nu_i^* h_i(x^*) \\ &\leq f(x^*) \end{aligned}$$

and therefore  $\lambda_i^* g_i(x^*) = 0 \quad \forall i$  □

## Sufficiency

Theorem | If the primal problem is convex (i.e.,  $f, g_i$  are convex functions,  $h_i$  are affine), and  $\tilde{x}, \tilde{\lambda}, \tilde{\nu}$  satisfy the KKT conditions, then  $\tilde{x}$  is primal optimal,  $(\tilde{\lambda}, \tilde{\nu})$  is dual optimal, and the duality gap is zero.

Proof: (2), (3)  $\Rightarrow \tilde{x}$  is feasible

(4)  $\Rightarrow L(x, \tilde{\lambda}, \tilde{\nu})$  is convex in  $x$

(1)  $\Rightarrow \tilde{x}$  is a minimizer of  $L(x, \tilde{\lambda}, \tilde{\nu})$ . Then

$$L_D(\tilde{\lambda}, \tilde{\nu}) = L(\tilde{x}, \tilde{\lambda}, \tilde{\nu})$$

$$= f(\tilde{x}) + \sum \tilde{\lambda}_i g_i(\tilde{x}) + \sum \tilde{\nu}_i h_i(\tilde{x})$$

$$= f(\tilde{x}) \quad \underbrace{\hspace{10em}}_{= 0 \text{ by (5)}} \quad \square$$

### Conclusion

If a constrained optimization problem is differentiable, convex, and a constraint qualification holds, then the KKT conditions are necessary and sufficient for primal/dual optimality (with zero duality gap). Thus, the KKT conditions can be used to solve such problems.

**Key** A. feasible ; Lagrangian ; Lagrange multipliers  
or dual variables

B. dual ; primal

C. duality gap ; convex ; affine

**Reference** Convex Optimization, Ch 5  
Boyd + Vandenberghe  
(available online)