

Entropic Graphs

Alfred O. Hero

Dept. EECS, Dept Biomed. Eng., Dept. Statistics

University of Michigan - Ann Arbor


`hero@eecs.umich.edu`

`http://www.eecs.umich.edu/~hero`

Collaborators: A. Almal, J. Costa, H. Neemuchwala, B. Ma, O. Michel

- Pattern Matching and Image Retrieval
- Entropic Euclidean Graphs
- A variant: K -point entropic graphs
- Application to US image registration
- Application to shape manifolds

Image Retrieval



QUERY




Figure 1: Yale face database <http://cvc.yale.edu/projects/yalefaces/yalefaces.html>

DATABASE


Multi-modality face Retrieval

Database of Visible/IR images



<http://www.equinoxsensors.com/products/HID.html>

SAR Target Classification



QUERY





Figure 2: SLICY canonical target images (54×54) at elevation 39° and different azimuth angles.

DATABASE

SAR Target Detection






Figure 3: SAR clutter image, target on boundary at column 305.


Image Registration



(a) Image I_1




(b) Image I_0




(c) Registration result

Figure 4: A multirate image registration example


Clustering Features in Feature Space




Multiple Mode Low Entropy Feature Density




Single Mode Low Entropy Feature Density



High Entropy Feature Density



Common Processing System



Objective: For given fitness criterion Q , find operator T which minimizes/maximizes Q

Our focus: entropic fitness criterion $Q(f)$

f : feature density over $x \in \mathbf{R}^d$

Some Popular Entropic Q 's

1. Shannon Entropy of feature density f

$$Q(f) = H(f) = - \int f(x) \ln f(x) dx$$

2. KL Divergence between feature densities f, g

$$Q(f, g) = D(f \| g) = \int f(x) \ln \left(\frac{f(x)}{g(x)} \right) dx$$

3. Jensen difference between feature densities f, g :

$$Q(f, g) = H(\epsilon f + (1 - \epsilon)g) - \epsilon H(f) - (1 - \epsilon)H(g)$$

4. Mutual information within joint feature density $f_{X,Y}$

$$Q(f_{X,Y}) = \text{MI}(X, Y) = \int \int f_{X,Y}(x, y) \ln \left(\frac{f_{X,Y}(x, y)}{f_X(x) f_Y(y)} \right) dx$$



Issue: How to estimate entropic Q from measured data?

Some possibilities:

1. Assume parameteric models for $f, g, f_{X,Y}$
2. Quantize feature space and use histograms
3. Non-parameteric density estimation of $f, g, f_{X,Y}$

Our Strategy: construct “entropic graphs” on features

A Set of Feature Samples and a Euclidean Spanning Graph




Minimal Euclidean Graphs: MST

Let $T_n = T(X_n)$ denote the possible sets of edges in the class of acyclic graphs spanning X_n (spanning trees).

The Euclidean Power Weighted MST achieves

$$L_\gamma^{\text{MST}}(X_n) = \min_{T_n} \sum_{e \in T_n} \|e\|^\gamma.$$




Minimal Euclidean graphs: k -NNG

Let $N_{k,i}(X_n)$ denote the possible sets of k edges connecting point x_i to all other points in X_n .

The Euclidean Power Weighted k -NNG is

$$L_\gamma^{k-NNG}(X_n) = \sum_{i=1}^n \min_{N_{k,i}(X_n)} \sum_{e \in N_{k,i}(X_n)} |e|^\gamma$$



MST for Two Different Samples




Figure 5:

Large n behavior of MST




Figure: MST and log MST weights as function of the number of samples.

Asymptotics: the BHH Theorem

Define the MST length functional

$$L_\gamma(X_n) = \min_{T_n} \sum_{e \in T_n} \|e\|^\gamma.$$

Theorem 1 [*Beardwood, Halton&Hammersley:1959*] Let $X_n = \{X_1, \dots, X_n\}$ be an i.i.d. realization from a Lebesgue density f with support $S \subset [0, 1]^d$.

$$\lim_{n \rightarrow \infty} L_\gamma(X_n) / n^{(d-\gamma)/d} = \beta_{L_\gamma, d} \int_S f(x)^{(d-\gamma)/d} dx, \quad (a.s.)$$

Or, letting $\alpha = (d - \gamma) / d$

$$\frac{1}{1 - \alpha} \ln L_\gamma(X_n) / n^\alpha \rightarrow H_\alpha(f) + c \quad (a.s.)$$

Rényi Entropy and Divergence

- Rényi Entropy of order α [Rényi:61,70]

$$H_{\alpha}(f) = \frac{1}{1-\alpha} \ln \int_S f^{\alpha}(x) dx$$

- Rényi α -divergence of fractional order $\alpha \in [0, 1]$

$$\begin{aligned} D_{\alpha}(f_1 \parallel f_0) &= \frac{1}{\alpha-1} \ln \int_S f_0 \left(\frac{f_1}{f_0} \right)^{\alpha} dx \\ &= \frac{1}{\alpha-1} \ln \int_S f_1^{\alpha} f_0^{1-\alpha} dx \end{aligned}$$

- α -Divergence vs. Kullback-Liebler divergence

$$\lim_{\alpha \rightarrow 1} D_{\alpha}(f_1 \parallel f_0) = \int f_1 \ln \frac{f_1}{f_0} dx.$$

Clustering via K-MST


Assume f is a mixture density of the form

$$f = (1 - \epsilon)f_1 + \epsilon f_o,$$


where

- f_o is a known (uniform) outlier density
- f_1 is an unknown target density
- $\epsilon \in [0, 1]$ is unknown mixture parameter


Annulus Feature Density



Uniform Outlier Density



Mixture Density



K -point Minimal Spanning Tree (K -MST)




Figure 6: *Clustering an annulus density from uniform noise via k -MST.*

K-MST Stopping Rule




Figure 7: *Left: k-MST curve for 2D annulus density with addition of uniform “outliers” has a knee in the vicinity of $n - k = 35$.*

Greedy partitioning approximation to K-MST




Figure 8: A smallest subset B_k^m is the union of the two cross hatched cells shown for the case of $m = 5$ and $k = 17$.

Extended BHH Theorem for Greedy K-MST

Fix $\rho \in [0, 1]$. If $k/n \rightarrow \rho$ then the length of the greedy partitioning K -MST satisfies [Hero&Michel:IT99]

$$L_\gamma(X_{n,k}^*) / (\lfloor \rho n \rfloor)^\alpha \rightarrow \beta_{L_\gamma, d} \min_{A: \int_A f \geq \rho} \int_S f^\alpha(x|x \in A) dx \quad (a.s.)$$

or, alternatively, with

$$H_\alpha(f|x \in A) = \frac{1}{1-\alpha} \ln \int_S f^\alpha(x|x \in A) dx$$

$$\frac{1}{1-\alpha} \ln L_\gamma(X_{n,k}^*) / (\lfloor \rho n \rfloor)^\alpha \rightarrow \beta_{L_\gamma, d} \min_{A: \int_A f \geq \rho} H_\alpha(f|x \in A) \quad (a.s.)$$




Figure 9: *Waterpouring construction of minimum entropy density.*

k-MST Influence Function




Figure 10: *MST and k-MST influence curves for Gaussian density on the plane.*

Extension of BHH to Divergence Estimation?

Question: How to go from

$$\frac{1}{1-\alpha} \ln \int f^\alpha(x) dx \quad \text{to} \quad \frac{1}{\alpha-1} \ln \int f^\alpha(x) g^{1-\alpha}(x) dx \quad ?$$

- $g(x)$: a reference density on \mathbf{R}^d
- Assume $f \ll g$, i.e. for all x such that $g(x) = 0$ we have $f(x) = 0$.
- Make measure transformation $M(x)$ such that $dx \rightarrow g(x)dx$ on $[0, 1]^d$.
Then for $Y_n = M(X_n)$

$$L_\gamma(Y_n)/n^\alpha \rightarrow \beta_{L_\gamma, d} \int \left(\frac{f(x)}{g(x)} \right)^\alpha g(x) dx, \quad (a.s.)$$




Figure 11: Top Left: i.i.d. sample from triangular distribution, Top Right: exact transformation, Bottom: after application of exact and empirical transformations.

Clustering Example

X_n is a sample from the mixture

$$f(x) = (1 - \varepsilon)g(x) + \varepsilon h(x)$$

$h(x)$ is uniform density on $[0, 1]^2$

$g(x)$ is triangular density on $[0, 1]^2$

ε is unknown

Objective: Detect deviation of f from triangular and cluster the uniform variates in the sample

Illustration




Figure 12: Left: A sample from triangle-uniform mixture density with $\varepsilon = 0.9$ in the transformed domain Y_n . Right: ROC curves of thresholded α -divergence test for deviation from g . Curves are decreasing in ε over the range $\varepsilon \in \{0.1, 0.3, 0.5, 0.7, 0.9\}$

Clustering Illustration





Figure 13: *Left: the K-MST implemented on the transformed scatterplot Y_n with $k = 230$. Right: same K-MST displayed in the original data domain.*

Bounds on Minimax Convergence Rate

Theorem 2 (Hero,Costa&Ma 2001) *Let $d \geq 2$ and $1 \leq \gamma \leq d - 1$.*

Assume X_1, \dots, X_n are i.i.d. random vectors over $[0, 1]^d$ with density $f \in \Sigma_d(\beta, l)$, $\beta, l > 0$, having support $S \subset [0, 1]^d$. Assume also that $f^{\frac{1}{2} - \frac{\gamma}{d}}$ is integrable. Then,

$$\begin{aligned} O\left(n^{-r_1(d, \beta)}\right) &\leq \\ \sup_{f \in \Sigma_d(\beta, l)} E \left[\left| L_\gamma(X_1, \dots, X_n) / n^{(d-\gamma)/d} - \beta_{L_\gamma, d} \int_S f^{(d-\gamma)/d}(x) dx \right|^p \right]^{1/p} \\ &\leq O\left(n^{-r_2(d, \gamma)}\right), \end{aligned}$$

where

$$r_1(d, \beta) = \min\left\{\frac{4\beta}{4\beta + d}, 1/2\right\} \quad r_2(d, \gamma) = \frac{\alpha}{\alpha + 1} \frac{1}{d}$$

and $\alpha = \frac{d-\gamma}{d}$.

Extension to Partition Approximations

$$L_{\gamma}^m(X_n) = \sum_{i=1}^{m^d} L_{\gamma}(X_n \cap Q_i) + b(m),$$




Figure 14: *Partition approximation.*

Theorem 3 (Hero, Costa & Ma 2001) *Let $L_\gamma^m(X_n)$ be a partition approximation to $L_\gamma(X_n)$. Under the same hypotheses as in the previous proposition, if $b(m) = O(m^{d-\gamma})$*

$$O\left(n^{-r_1(d,\beta)}\right) \leq \sup_{f \in \Sigma_d(\beta, l)} E \left[\left| L_\gamma^{m(n)}(X_1, \dots, X_n) / n^{(d-\gamma)/d} - \beta_{L_\gamma, d} \int_S f^{(d-\gamma)/d}(x) dx \right|^p \right]^{1/p} \leq O\left(n^{-r_3(d, \gamma)}\right),$$

where

$$r_3(d, \gamma) = \frac{\alpha}{\frac{d-1}{\gamma} \alpha + 1} \frac{1}{d}.$$

This bound is attained by choosing the progressive-resolution sequence $m = m(n) = n^{1/[d(\frac{d-1}{\gamma} \alpha + 1)]}$.

Application: Registration of Breast Images




Figure 15: Three ultrasound breast scans. From top to bottom are: case 151, case 142 and case 162.

MI Registration of Gray Levels (Viola&Wells:ICCV95)

- X : a $N \times N$ image (lexicographically ordered)
- $X(k)$: image gray level at pixel location k
- X_0 and X_1 : primary and secondary images to be registered


Hypothesis: $\{(X_0(k), X_i(k))\}_{k=1}^{N^2}$ are i.i.d. r.v.'s with j.p.d.f

$$f_{0,i}(x_0, x_1), \quad x_0, x_1 \in \{0, 1, \dots, 255\}$$


Mutual Information (MI) criterion: $T = \operatorname{argmax}_{T_i} \hat{\text{MI}}$

where $\hat{\text{MI}}$ is an estimate of

$$\text{MI}(f_{0,i}) = \int \int f_{0,i}(x_0, x_1) \ln f_{0,i}(x_0, x_1) / (f_0(x_0) f_i(x_1)) dx_1 dx_0. \quad (1)$$



(a) Image I^R



(b) Image I^T

Figure 16: Single Pixel Coincidences (Left and right: reference image I^R at 0° and rotated image I^T at 8°)

Single-Pixel Scatterplot $(Z_j^R, Z_j^T)_{j=1}^p$





Figure 17: Grey level scatterplots. 1st Col: target=reference slice. 2nd Col: target = reference+1 slice.

Higher Level Features


Disadvantages of single-pixel features:

- Only depends on histogram of single pixel pairs
- Insensitive to spatial reordering of pixels in each image
- Difficult to select out grey level anomalies (shadows, speckle)
- Spatial discriminants fall outside of single pixel domain
- **Alternative:** Aggregate spatial features

Local Tags



(a) Image I^R



(b) Image I^T

Figure 18: Local Tag Coincidences

Generalization: α -MI Registration of Coincident Features

- X : a $N \times N$ US image (lexicographically ordered)
- $Z = Z(X)$: a general image feature vector in a d -dimensional feature space

Let $\{Z_0(k)\}_{k=1}^K$ and $\{Z_i(k)\}_{k=1}^K$ be features extracted from X_0 and X_i at K pairs of identical spatial locations

α -MI coincident-feature criterion

$$T = \operatorname{argmax}_{T_i} \hat{M}I_{\alpha}$$

where $\hat{M}I_{\alpha}$ is an estimate of

$$MI_{\alpha}(f_{0,i}) = \frac{1}{\alpha - 1} \log \int \int f_{0,i}^{\alpha}(z_0, z_1) f_0^{1-\alpha}(z_0) f_i^{1-\alpha}(z_1) dz_1 dz_0. \quad (2)$$

α -MI and Decision Theoretic Error Exponents

H_0 : $Z_0(k), Z_i(k)$ independent

H_1 : $Z_0(k), Z_i(k)$ o.w.

Bayes probability of error

$$P_e(n) = \beta(n)P(H_1) + \alpha(n)P(H_0)$$

Chernoff bound

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \log P_e(n) = - \sup_{\alpha \in [0,1]} \{ (1 - \alpha) \text{MI}_\alpha(f_{0,i}) \}.$$

ICA Features

Decomposition of $M \times M$ tag images $Y(k)$ acquired at $k = 1, \dots, K$ spatial locations

$$Y(k) = \sum_{p=1}^P a_{kp} S_p$$

- $\{S_k\}_{k=1}^P$: statistically independent components
- a_{kp} : projection coefficients of tag $Y(k)$ onto component S_p
- $\{S_k\}_{k=1}^P$ and P : selected via FastICA
- Feature vector for coincidence processing:

$$Z(k) = [a_{k1}, \dots, a_{kP}]^T$$

ICA feature basis for US breast images




Figure 19: *Estimated ICA basis set using FastICA*

Simpler Objective Function: α -Jensen Difference

1. Extract features from reference and transformed target images:

$$X_m = \{X_i\}_{i=1}^m \quad \text{and} \quad Y_n = \{Y_i\}_{i=1}^n$$

2. Construct following MST function on X_m and Y_n


$$\Delta L = \ln L_\gamma(X_m \cup Y_n) / (n+m)^\alpha - \frac{m}{n+m} \ln L_\gamma(X_m) / m^\alpha - \frac{n}{n+m} \ln L_\gamma(Y_n) / n^\alpha$$

3. Minimize ΔL_γ over transformations producing Y_n .


$$(1 - \alpha)^{-1} \Delta L \rightarrow H_\alpha(\epsilon f_x + (1 - \epsilon) f_y) - \epsilon H_\alpha(f_x) - (1 - \epsilon) H_\alpha(f_y)$$

where $\epsilon = \frac{m}{m+n}$

Illustration




(a)




(b)

Figure 20: MST demonstration for misaligned images



(a)



(b)

Figure 21: MST for aligned images. “x” denotes reference while “o” denotes a candidate image in the DEM database.

Quantitative Performance Comparisons for US Registration





Figure 22: *US registration MSE comparisons.*

Application to Classification of Shape Manifolds



Intrinsic Dimension of Shape Manifolds

Focus applications: face compression and recognition, lung nodule classification, minimal feature selection

Property: $N \times N$ images $\{X_i\}_{i=1}^n$ of smooth shapes lie on a proper d -dimensional subspace of \mathbf{R}^{N^2} :

$$X_i \in \mathcal{S} \subset \mathbf{R}^{N^2}$$

Implications:

1. *Extrinsic dimension $N^2 > \text{Intrinsic dimension } d$*
2. N^2 -dimensional image density $f(x)$ is concentrated on \mathcal{S} and has *extrinsic entropy*

$$\int_{\mathbf{R}^{N^2}} f^\alpha(x) dx = 0$$

MST and Intrinsic Dimension

Proposition 1 *Let the manifold $S \subset \mathbf{R}^{N^2}$ have dimension d_o . Then if $X \in \mathbf{R}^{N^2}$ are i.i.d. realizations from a Lebesgue density $f(x)$ on S :*

$$E[L_\gamma(X_1, \dots, X_n)]/n^{(d-\gamma)/d} \rightarrow \begin{cases} \infty, & d < d_o \\ \beta_{L_\gamma, d_o} \int_S f^{(d_o-\gamma)/d_o}(x) dx, & d = d_o \\ 0, & d > d_o \end{cases}$$

\Rightarrow **Estimation:** of $[d, H_\alpha]$ via LS solution $[\hat{\alpha}, \hat{C}]$ to

$$\ln E[L_\gamma(X_1, \dots, X_n)] = \overbrace{(d-\gamma)/d}^{\alpha} \ln n + C + \epsilon_n, \quad n \geq n_o \dots$$

- $\hat{d} = \gamma/(1 - \hat{\alpha})$
- $\hat{H}_\alpha = \hat{C}/(1 - \hat{\alpha})$

Estimation Procedure Applied to Shape Database

Intrinsic dimensionality of shape manifold is estimated using MST:

Initialize: Using entire database of shapes construct distance matrix E on face manifold via *isomap* (Tenenbaum *etal* 2000).

For $n = n_o, \dots, n_1$

Repeat

Randomly select n face images from database

Construct MST over n face images using E matrix
end repeat

Compute sample average MST length $\hat{L}_\gamma(n)$

end for

Estimate d and H_α from $\{\hat{L}_\gamma(n)\}_n$ via LS

Application to Yale Face Database



Figure 23: *Samples of a face from Yale database.*

- Faces of 4 persons investigated
- 585 poses and illumination conditions for each person
- extrinsic dimension = 4096, $\gamma = 1$

Application to Yale Face Database




Figure 24: Avg. *intrinsic* MST length for three faces.

Application to Yale Face Database: Results

Single Faces

\Rightarrow Dimension of single face manifold: $3 < \hat{d} < 4$

\Rightarrow Entropy of single face manifold: $28 < \hat{H}_\alpha < 32$

Multiple Faces

\Rightarrow Dimension of any 3 face manifold: $6 < \hat{d} < 7$

\Rightarrow Entropy of 3 face manifold: $34 < \hat{H}_\alpha < 40$

Conclusions

1. Entropic graphs can be used to estimate α -entropy and α -divergence
2. MST and k-NN applied to high dimensional feature-based image registration
3. Clustering using entropic K -point graphs
4. Extensions to larger class of continuous quasi-additive graphs (Yukich)
5. Can use entropic graphs to explore multivariate shape distributions
6. Can also handle case of unknown shape manifold dimension