Imaging Applications of Stochastic Minimal Graphs

A. O. Hero*, B. Ma †, O. Michel #

University of Michigan*, Intervideo, Inc.†, University of Nice#

hero@eecs.umich.edu

http://www.eecs.umich.edu/~hero

Outline

1. Divergence and entropy measures
2. Plug-in and Graph Estimators of Entropy
3. Theoretical comparisons
4. Applications



Figure 1: SAR clutter image



(a) 142° (b) 147° (c) 152° (d) 157° (e) 163°



(f) 169° (g) 175° (h) 187° (i) 193°

Figure 2: SLICY canonical target images.

Statistical Framework

- X: an image
- $Z = Z(X)$: an image feature vector
- Θ: a parameter space
- $f(z|\theta)$: feature density (likelihood)
- X_R a reference image
- $\{X_i\}_i$ a database of K images

$$Z^R = Z(X_R) \quad \sim \quad f(z|\theta_R)$$

$$Z^i = Z(X_i) \quad \sim \quad f(z|\theta_i), \quad i = 1\ldots,K$$

$\Rightarrow$ Similarity btwn X_i, X_R lies in similarity btwn models

Divergence Measures

Refs: [Csiszár:67,Basseville:SP89]

Define densities

$$f_i = f(z|\theta_i), \quad f_R = f(z|\theta_R)$$

The Rényi α-divergence of fractional order $\alpha \in [0,1]$ [Rényi:61,70]

$$\begin{aligned} D_\alpha(f_i \parallel f_R) = D(\theta_i \| \theta_R) &= \frac{1}{\alpha - 1} \ln \int f_R \left(\frac{f_i}{f_R} \right)^\alpha dx \\ &= \frac{1}{\alpha - 1} \ln \int f_i^\alpha f_R^{1-\alpha} dx \end{aligned}$$

Rényi α-Divergence: Special cases

- α-Divergence vs. Hellinger Affinity

$$D_{\frac{1}{2}}(f_i \parallel f_R) = \ln \left(\int \sqrt{f_i f_R} dx \right)^2$$

- α-Divergence vs. Kullback-Liebler divergence

$$\lim_{\alpha \to 1} D_\alpha(f_i, f_R) = \int f_R \ln \frac{f_R}{f_i} dx.$$

- α-Divergence vs. α-MI

$$\mathrm{MI}_\alpha(f_{X,Y}) = D_\alpha(f_{X,Y} \| f_X f_Y)$$

- α-Divergence vs. α-Entropy

$$H_\alpha(f_i) = D_\alpha(f_i \| U[0,1]) = \frac{1}{1-\alpha} \ln \int f_i^\alpha(x) dx$$

Indexing via α-divergence

Clairvoyant indexing rule:

$$X_i \prec X_j \quad \Leftrightarrow \quad D_\alpha(f_i \| f_R) < D_\alpha(f_j \| f_R)$$

Indexing problem: find θ_i attaining $\min_{\theta_i \neq \Theta_R} D_\alpha(\theta_i \| \theta_R)$

1. Image classification: f_i index model classes [Stoica&etal:INRIA98]
2. Target detection: f_R is noise reference and f_i are target references. Declare detection if $\min_{\theta_i \neq \Theta_R} D_\alpha(\theta_i \| \theta_R) >$ threshold

Methods of Divergence Estimation

- $Z = Z(X)$: a statistic (MI, reduced rank feature, etc)
- $\{Z^{(n)}\}$: n i.i.d. realizations from $f(Z;\theta)$

Objective: Estimate $\hat{D}_\alpha(f_i \| f_R)$ from $\{Z_i^{(n)}, Z_R^{(n)}\}$'s

1. Parametric density estimation methods
2. Non-parametric density estimation methods
3. Non-parametric minimal-graph estimation methods

Non-parametric estimation methods

Given i.i.d. sample $X = \{X_1, \ldots, X_n\}$

Density "plug-in" estimator

$$H_\alpha(\hat{f}_n) = \frac{1}{1-\alpha} \ln \int_{\mathbf{R}^d} \hat{f}^\alpha(x) dx$$

Previous work limited to Shannon entropy $H(f) = -\int f(x) \ln f(x) dx$

- Histogram plug-in [Gyorfi&VanDerMeulen:CSDA87]
- Kernel density plug-in [Ahmad&Lin:IT76]
- Sample-spacing plug-in [Hall:JMS86] $(d = 1)$
  - Performance degrades as density f becomes non smooth
  - Unclear how to robustify $\hat{f}$ against outliers
  - d-dimensional integration might be difficult
  - $\Rightarrow$ function $\{f(x) : x \in \mathbf{R}^d\}$ over-parameterizes entropy functional

Direct α-entropy estimation

- MST estimator of α-entropy [Hero&Michel:IT99]:

$$\hat{H}_\alpha = \frac{1}{1-\alpha} \ln L_\gamma(\mathcal{X}_n)/n^{-\alpha}$$

  - Direct entropy estimator: faster convergence for nonsmooth densities
  - Parameter α is varied by varying interpoint distance measure
  - Optimally pruned k-MST graphs robustify $\hat{f}$ against outliers
  - Greedy multi-scale MST approximations reduce combinatorial complexity

Minimal Graphs: Minimal Spanning Tree (MST)

Let $\mathcal{M}_n = \mathcal{M}(\mathcal{X}_n)$ denote the possible sets of edges in the class of acyclic graphs spanning $\mathcal{X}_n$ (spanning trees).

The Euclidean Power Weighted MST achieves

$$L_{\mathrm{MST}}(\mathcal{X}_n) = \min_{\mathcal{M}_n} \sum_{e \in \mathcal{M}_n} \|e\|^{\gamma}.$$



Figure 3: *A sample data set and the MST*

Minimal Graphs: Pruned MST

Fix k, $1 \leq k \leq n$.

Let $\mathcal{M}_{n,k} = \mathcal{M}(x_{i_1}, \ldots, x_{i_k})$ be a minimal graph connecting k distinct vertices $x_{i_1}, \ldots, x_{i_k}$.

The k-MST $\mathrm{T}^*_{n,k} = \mathrm{T}^*(x_{i_1^*}, \ldots, x_{i_k^*})$ is minimum of all k-point MST's

$$L^*_{n,k} \quad = \quad L^*(\mathcal{X}_{n,k}) = \min_{i_1,\ldots,i_k} \min_{\mathcal{M}_{n,k}} \sum_{e \in \mathcal{M}_{n,k}} \|e\|^\gamma$$



Figure 4: *k-MST for 2D torus density with and without the addition of uniform "outliers".*

Convergence of MST



Figure 5: *2D Triangular vs. Uniform sample study for MST.*



Figure 6: *MST and log MST weights as function of number of samples for 2D uniform vs. triangular study.*



Figure 7: *Continuous quasi-additive euclidean functional satisfies "self-similarity" property on any scale.*

Asymptotics: Plug-in estimation of $H_\alpha(f)$

Class of Hölder continuous functions over $[0,1]^d$

$$\Sigma_d(\kappa,c) = \left\{ f(x) : |f(x) - p_x^{\lfloor \kappa \rfloor}(z)| \leq c \, \|x-z\|^\kappa \right\}$$

Class of functions of Bounded Variation (BV) over $[0,1]^d$

$$\mathrm{BV}_d(c) = \left\{ f(x) : \sup_{\{x_i\}} \sum_i |f(x_i) - f(x_{i-1})| \leq c \right\}.$$

Proposition 1 (Hero&Ma:IT01) *Assume that $f^\alpha \in \Sigma_d(\kappa,c)$. Then, if $\hat{f}^\alpha$ is a* **minimax** *estimator*

$$\sup_{f^\alpha \in \Sigma_d(\kappa,c)} E^{1/p}\left[\left| \int \widehat{f}^\alpha(x)dx - \int f^\alpha(x)dx \right|^p \right] = O\left(n^{-\kappa/(2\kappa+d)} \right)$$

Asymptotics: Minimal-graph estimation of $H_\alpha(f)$

Proposition 2 (Hero&Ma:IT01) *Let $d \geq 2$ and $\alpha = (d-\gamma)/d \in [1/2,(d-1)/d]$. Assume that $f^\alpha \in \Sigma_d(\kappa,c)$ where $\kappa \geq 1$ and $c < \infty$. Then for any continuous quasi-additive Euclidean functional L_γ*

$$\sup_{f^\alpha \in \Sigma_d(\kappa,c)} E^{1/p}\left[\left|\frac{L_\gamma(X_1,\ldots,X_n)}{n^\alpha} - \beta_{L_\gamma,d}\int f^\alpha(x)dx\right|^p\right] \leq O\left(n^{-1/(d+1)}\right)$$

Conclude: minimal-graph estimator converges faster for

$$\kappa < \frac{d}{d-1}$$

As $\Sigma_d(1,c) \subset \mathrm{BV}_d(c)$, we have

Corollary 1 (Hero&Ma:IT01) *Let $d \geq 2$ and $\alpha = (d-\gamma)/d \in [1/2,(d-1)/d]$. Assume that f^α is of bounded variation over $[0,1]^d$. Then*

$$\sup_{f^\alpha \in \mathrm{BV}_d(c)} E^{1/p}\left[\left|\int \widehat{f}^\alpha(x)dx - \beta_{L_\gamma,d}\int f^\alpha(x)dx\right|^p\right] \geq O\left(n^{-1/(d+2)}\right)$$

$$\sup_{f^\alpha \in \mathrm{BV}_d(c)} E^{1/p}\left[\left|\frac{L_\gamma(X_1,\ldots,X_n)}{n^\alpha} - \beta_{L_\gamma,d}\int f^\alpha(x)dx\right|^p\right] \leq O\left(n^{-1/(d+1)}\right)$$

Observations

- Minimal graph rates valid for MST, k-NN graph, TSP, Steiner Tree, etc
- Analogous rate bound holds for progressive-resolution algorithm

$$L_\gamma^G(\mathcal{X}_n) = \sum_{i=1}^{m^d} L_\gamma(\mathcal{X}_n \cap Q_i)$$

  $\{Q_i\}$ is uniform partition of $[0,1]^d$ into cell volumes $1/m^d$

- Optimal sequence of cell volumes is:

$$m^{-d} = n^{-1/(d+1)}$$

- These results also apply to greedy multi-resolution k-MST

Application: Image Registration

Two independent data samples from unknown distributions

- $X = [X_1, \ldots, X_m] \sim f(x)$
- $Y = [Y_1, \ldots, Y_n] \sim g(x)$

Suppose: $g(x) = f(Ax+b)$, $A^T A = I$

Objective: find rigid transformation A, b

- Two methods:

1. α-MI of $\{(X_i, Y_i)\}_{i=1}^n$
2. α-Entropy of $\{X_i\}_{i=1}^m \cup \{Y_i\}_{i=1}^n$



(a) (b) (c)

Figure 8: Reference and target SAR/DEM images

$O(n^{-1/(2d+1)})$ algorithm for α-MI estimation

$$\mathrm{MI}_\alpha(X,Y) = \frac{1}{\alpha-1}\ln\int f_{X,Y}^\alpha(x,y)(f_X(x)f_Y(y))^{1-\alpha}dxdy.$$

Algorithm:

1. `Kernel estimates` $\hat{f}_X, \hat{f}_Y$ $(O(n^{-1/(d+2)}))$
2. `Uniformizing probability transformations:` $\tilde{X} = F_X(X),\ \tilde{Y} = F_Y(Y)$
3. `Graph entropy estimate of` $\mathrm{MI}_\alpha(X,Y)$ $(O(n^{-1/(2d+1)}))$

$$\frac{L_\gamma(\{(\tilde{X}_1,\tilde{Y}_1),\ldots,(\tilde{X}_n,\tilde{Y}_n)\})}{n^\alpha} \quad \to \quad \beta_{L_\gamma,d}\int f_{\tilde{X},\tilde{Y}}^\alpha(x,y)dxdy$$

$$= \quad \beta_{L_\gamma,d}\int f_{X,Y}^\alpha(x,y)(f_X(x)f_Y(y))^{1-\alpha}dxdy \quad (w.p$$

$O(n^{-1/(d+1)})$ criterion: α-Jensen difference

- Jensen's difference btwn f_0, f_1:

$$\Delta J_\alpha = H_\alpha(\varepsilon f_1 + (1-\varepsilon) f_0) - \varepsilon H_\alpha(f_1) - (1-\varepsilon) H_\alpha(f_0) \geq 0$$

- f_0, f_1 are two densities, ε satisfies $0 \leq \varepsilon \leq 1$
- Let X, Y be i.i.d. features extracted from two images

$$X = \{X_1, \ldots, X_m\}, \quad Y = \{Y_1, \ldots, Y_n\}$$

- Each realization in *unordered* sample $Z = \{X, Y\}$ has marginal

$$f_Z(z) = \varepsilon f_X(z) + (1-\varepsilon) f_Y(z), \quad \varepsilon = \frac{m}{n+m}$$

- α-Jensen difference for rigid transformation T

$$\Delta J_\alpha(\mathrm{T}) = H_\alpha(\varepsilon f_X + (1-\varepsilon) f_Y) - \underbrace{\varepsilon H_\alpha(f_X) - (1-\varepsilon) H_\alpha(f_Y)}_{constant}$$



Figure 9: Reference and target SAR/DEM images



(a) (b)

Figure 10: MST demonstration for misaligned images



(a) (b)

Figure 11: MST demonstration for aligned images

Conclusions

1. α-divergence for indexing can be justified via decision theory
2. Non-parametric estimation of Jensen's difference is low complexity alternative to α-divergence estimation
3. Non-parametric estimation of Jensen's difference is possible without density estimation
4. Minimal-graph estimation outperforms plug-in estimation for non-smooth densities